



*Società
Medico Chirurgica
di Ferrara*

TRATTAMENTO MININVASIVO

PECTUS EXCAVATUM

Claudio Vella

SC Chirurgia Pediatrica

Azienda Ospedaliero Universitaria Ferrara



**Università
degli Studi
di Ferrara**



**SERVIZIO SANITARIO REGIONALE
EMILIA-ROMAGNA
Azienda Ospedaliero - Universitaria di Ferrara
Azienda Unità Sanitaria Locale di Ferrara**

A 10-Year Review of a Minimally Invasive Technique for the Correction of Pectus Excavatum

By Donald Nuss, Robert E. Kelly, Jr, Daniel P. Croitoru, and Michael E. Katz
Norfolk, Virginia

Purpose: The aim of this study was to assess the results of a 10-year experience with a minimally invasive operation that requires neither cartilage incision nor resection for correction of pectus excavatum.

Methods: From 1987 to 1996, 148 patients were evaluated for chest wall deformity. Fifty of 127 patients suffering from pectus excavatum were selected for surgical correction. Eight older patients underwent the Ravitch procedure, and 42 patients under age 15 were treated by the minimally invasive technique. A convex steel bar is inserted under the sternum through small bilateral thoracic incisions. The steel bar is inserted with the convexity facing posteriorly, and when it is in position, the bar is turned over, thereby correcting the deformity. After 2 years, when permanent remodeling has occurred, the bar is removed in an outpatient procedure.

Results: Of 42 patients who had the minimally invasive procedure, 30 have undergone bar removal. Initial excellent results were maintained in 22, good results in four, fair in two, and poor in two, with mean follow-up since surgery of 4.6 years (range, 1 to 9.2 years). Mean follow-up since bar removal is 2.8 years (range, 6 months to 7 years). Average blood loss was 15 mL. Average length of hospital stay was 4.3

days. Patients returned to full activity after 1 month. Complications were pneumothorax in four patients, requiring thoracostomy in one patient; superficial wound infection in one patient; and displacement of the steel bar requiring revision in two patients. The fair and poor results occurred early in the series because (1) the bar was too soft (three patients), (2) the sternum was too soft in one of the patients with Marfan's syndrome, and (3) in one patient with complex thoracic anomalies, the bar was removed too soon.

Conclusions: This minimally invasive technique, which requires neither cartilage incision nor resection, is effective. Since increasing the strength of the steel bar and inserting two bars where necessary, we have had excellent long-term results. The upper limits of age for this procedure require further evaluation.

J Pediatr Surg 33:545-552. Copyright © 1998 by W.B. Saunders Company.

INDEX WORDS: Pectus excavatum, minimally invasive surgery, computed tomography scans in chest disease, thorax abnormalities.









Fig 3. Age at time of surgery.

A 10-Year Review of a Minimally Invasive Technique for the Correction of Pectus Excavatum

By Donald Nuss, Robert E. Kelly, Jr, Daniel P. Croitoru, and Michael E. Katz
Norfolk, Virginia



Short Nuss bar procedure

Hans Kristian Pilegaard^{1,2}

¹Department of Cardiothoracic and Vascular Surgery, Aarhus University Hospital, Skejby, Aarhus, Denmark; ²Department of Clinical Medicine, Aarhus University, Aarhus, Denmark

Correspondence to: Hans Kristian Pilegaard. Department of Cardiothoracic and Vascular Surgery, Aarhus University Hospital, Skejby, Palle Juul-Jensens Boulevard 99, DK-8200 Aarhus N, Denmark. Email: pilegaard@dadlnet.dk.

The Nuss procedure is now the preferred operation for surgical correction of pectus excavatum (PE). It is a minimally invasive technique, whereby one to three curved metal bars are inserted behind the sternum in order to push it into a normal position. The bars are left in situ for three years and then removed. This procedure significantly improves quality of life and, in most cases, also improves cardiac performance. Previously, the modified Ravitch procedure was used with resection of cartilage and the use of posterior support. This article details the new modified Nuss procedure, which requires the use of shorter bars than specified by the original technique. This technique facilitates the operation as the bar may be guided manually through the chest wall and no additional stabilizing sutures are necessary.

Keywords: Pectus excavatum repair (PE repair); Nuss; short bar; minimally invasive surgery

Submitted Mar 14, 2016. Accepted for publication Jul 12, 2016.

doi: 10.21037/acs.2016.09.06

View this article at: <http://dx.doi.org/10.21037/acs.2016.09.06>



Figure 4 Bar in an oblique position.




Figure 5 Crossed bars.




Figure 6 Bars in an angle. Same exit on the left side.




Figure 7 The asymmetric location of the bar.


Mininvasivo = Torascopia

Modifiche ed innovazioni :

Lunghezza barre

Mezzi di fissazione

Numero di barre



Symmetric Pectus Excavatum

